

Magnus Force and Aharonov-Bohm Effect in Superfluids

Edouard Sonin

Racah Institute of Physics,
Hebrew University of Jerusalem
and

Ioffe Physical Technical Institute, Solid State Physics Division,
St. Petersburg, Russia

February 1, 2008

Abstract

The paper addresses the problem of the transverse force (Magnus force) on a vortex in a Galilean invariant quantum Bose liquid. Interaction of quasiparticles (phonons) with a vortex produces an additional transverse force (Iordanskii force). The Iordanskii force is related to the acoustic Aharonov–Bohm effect. Connection of the effective Magnus force with the Berry phase is also discussed.

1 Introduction

In classical hydrodynamics it has long been known that if the vortex moves with respect to a liquid there is a force on the vortex normal to the vortex velocity [1]. This is the Magnus force, which is a particular case of a force on a body immersed into a liquid with a flow circulation around it (the Kutta-Joukowski theorem). An example of it is the lift force on an airplane wing.

The key role of the Magnus force in vortex dynamics became clear from the very beginning of studying superfluid hydrodynamics. In the pioneer article on the subject Hall and Vinen [2] defined the superfluid Magnus force as a force between a vortex and a superfluid. Therefore it was proportional to the superfluid density ρ_s . But in the two-fluid hydrodynamics the superfluid Magnus force is not the only force on the vortex transverse to its velocity: there was also a transverse force between the vortex and quasiparticles moving with respect to the vortex. The transverse force from rotons was found by Lifshitz and Pitaevskii [3] from the quasiclassical scattering theory. Later Iordanskii [4] revealed the transverse force from phonons. The analysis done in Ref. [5] (see also Refs. [6, 7]) demonstrated that the Lifshitz–Pitaevskii force for rotons and the Iordanskii force for phonons originate from interference between quasiparticles which move past the vortex on the left and on the right sides with different phase shifts, like in the Aharonov–Bohm effect [8]. Since the phase shift depends on the circulation which is a topological charge for a vortex, this is a clear indication of connection between the transverse quasiparticle force and topology.

Later on the analogy between wave scattering by vortex and electron scattering by the magnetic-flux tube (the Aharonov–Bohm effect) was studied in classical hydrodynamics for water surface waves (the acoustic Aharonov–Bohm effect [9, 10]). Scattering of the light by a vortex also results in the Aharonov–Bohm interference (the optical Aharonov–Bohm effect [11]). As follows from Ref. [5], the Aharonov–Bohm interference always produces a transverse force on the vortex, or the fluxon. For the original Aharonov–Bohm effect this force is discussed by Shelankov [12].

The Magnus force on a vortex in a superconductor was introduced by Nozières and Vinen [13]. The total transverse force on a vortex is responsible for the Hall effect in the mixed state. In superconductors not only quasiparticles, but also impurities produce an additional transverse force on the vortex. A reader can find discussion of this problem in the review by Kopnin [14].

Despite a lot of work done to understand and calculate the Magnus force, it remained to be a controversial issue. Ao and Thouless [15] pointed out a connection of the Magnus force with the Berry phase [16] which is the phase variation of the quantum-mechanical wave function resulting from transport of the vortex round a close loop. From the Berry-phase analysis Ao and Thouless concluded that the effective Magnus force is proportional to the

superfluid density, and there is no transverse force on the vortex from quasi-particles and impurities [15, 17, 18]. Such conclusion disagreed with the previous calculations of the Magnus force in superfluids and superconductors, and therefore generated a vivid discussion [19, 20, 21].

The present paper addresses this problem [22] which is very important for many issues in modern condensed matter physics, field theory, and cosmology. The analysis is based on studying momentum balance in the area around a moving vortex without using any preliminary concept of a force. The word *force* is a label to describe a transfer of momentum between two objects. A careful approach is to give these labels to the various terms in the momentum balance equation only *after* derivation of this equation. I restrict myself with the problem of the Galilean invariant quantum Bose liquid. For a weakly nonideal Bose gas one can use the Gross–Pitaevskii theory [23]. In this theory the liquid is described by the nonlinear Schrödinger equation. At large scales the nonlinear Schrödinger equation yields the hydrodynamics of an ideal inviscous liquid. In presence of an ensemble of sound waves (phonons) with the Planck distribution, which is characterized by a locally defined normal velocity $\vec{v}_n$ (the drift velocity of the Planck distribution), one obtains two-fluid hydrodynamics. Eventually the problem of the vortex motion in presence of the phonon normal fluid is a problem of hydrodynamics.

The paper starts from discussion of the Magnus force in classical hydrodynamics (Sec. 2). In Sec. 3 I define the superfluid Magnus force and the force from quasiparticles scattered by a vortex. Connection between the Gross–Pitaevskii theory and the two-fluid hydrodynamics is discussed in Sec. 4. In Sec. 5 I discuss scattering of a sound wave by a vortex and show that the standard scattering theory fails to give a conclusive result on the transverse force because of small-scattering-angle divergence of the scattering amplitude. The next Sec. 6 gives a solution of the sound equation around the vortex, which is free from the small-angle divergence. Using this solution in the momentum balance one obtains the equation of vortex motion, which contains the Iordanskii force. The momentum transfer responsible for the Iordanskii force occurs at small scattering angles where a phenomenon analogous to the Aharonov–Bohm effect is important: an interference between the waves on the left and on the right from the vortex with different phase shifts after interaction with the vortex. Section 7 shows how to derive the transverse force from the exact partial-wave solution of the Aharonov–Bohm problem for electrons. In our analysis the force on the vortex originates from

scattering of noninteracting quasiparticles. This is a valid assumption since normally the phonon mean-free path essentially exceeds the scale where the force arises (the phonon wavelength) [24]. But in order to know the effect of this force on the whole superfluid, it is important to investigate how the force is transmitted to distances much larger than the mean-free path where the two-fluid hydrodynamics becomes valid. It is done in Sec. 8. The effect of the force at very large distances is also important for discussion of the Berry phase in Sec.9. The transverse force creates a circulation of the normal velocity at very large distances from an isolated vortex. Taking into account this circulation, the Berry phase yields a correct value of the transverse force, which agrees with the result derived from the momentum balance.

The present paper does not discuss experimental aspects of this problem, which are addressed in other reviews devoted to rotating ^4He and ^3He [6, 26, 27]).

2 The Magnus force in classical hydrodynamics

It is worth to remind how the Magnus force appears in classical hydrodynamics. Let us consider a cylinder immersed in an incompressible inviscous liquid. There is a potential circular flow around the cylinder with the velocity

$$\vec{v}_v(\vec{r}) = \frac{\vec{\kappa} \times \vec{r}}{2\pi r^2} . \quad (1)$$

Here $\vec{r}$ is the position vector in the plane xy , the axis z is the axis of the cylinder, and $\vec{\kappa}$ is the circulation vector along the axis z . In classical hydrodynamics the velocity circulation $\kappa = \oint \vec{v}_v \cdot d\vec{l}$ may have arbitrary values. There is also a fluid current past the cylinder with a transport velocity $\vec{v}_{tr}$ and the net velocity is

$$\vec{v}(\vec{r}) = \vec{v}_v(\vec{r}) + \vec{v}_{tr} . \quad (2)$$

This expression is valid at distances r much larger than the cylinder radius. At smaller distances one should take into account that the flow with the velocity $\vec{v}_{tr}$ cannot penetrate into the cylinder, but velocity corrections due to this effect decrease as $1/r^2$ and are not essential for the further analysis.

The Euler equation for the liquid is

$$\frac{\partial \vec{v}}{\partial t} + (\vec{v} \cdot \vec{\nabla}) \vec{v} = -\frac{1}{\rho} \vec{\nabla} P . \quad (3)$$

Here ρ is the liquid density and P is the pressure.

Assuming that the cylinder moves with the constant velocity $\vec{v}_L$, i.e., replacing the position vector $\vec{r}$ by $\vec{r} - \vec{v}_L t$, one obtains that

$$\frac{\partial \vec{v}}{\partial t} = -(\vec{v}_L \cdot \vec{\nabla}) \vec{v} . \quad (4)$$

Then the Euler equation (3) yields the Bernoulli law for the pressure:

$$P = P_0 - \frac{1}{2} \rho [\vec{v}(\vec{r}) - \vec{v}_L]^2 = P'_0 - \frac{1}{2} \rho \vec{v}_v(\vec{r})^2 - \rho \vec{v}_v(\vec{r}) \cdot (\vec{v}_{tr} - \vec{v}_L) . \quad (5)$$

Here P_0 and $P'_0 = P_0 - \frac{1}{2} \rho (\vec{v}_{tr} - \vec{v}_L)^2$ are constants, which are of no importance for the following derivation. Figure 1 shows that due to superposition of two fluid motions given by Eq. (2) the velocity above the cylinder is higher than below the cylinder. According to the Bernoulli law, the pressure is higher in the area where the velocity is lower. As a result of it, a liquid produces a force on the cylinder normal to the relative velocity of the liquid with respect to the cylinder. This is a lift force, or the Magnus force.

In order to find the whole force, we must consider the momentum balance for a cylindrical region of a radius r_0 around the cylinder (see Fig. 1). The momentum conservation law requires that the external force $\vec{F}$ on the cylinder is equal to the momentum flux through the entire cylindrical boundary in the reference frame moving with the vortex velocity $\vec{v}_L$. The momentum-flux tensor is

$$\Pi_{ij} = P \delta_{ij} + \rho v_i(\vec{r}) v_j(\vec{r}) , \quad (6)$$

or in the reference frame moving with the vortex velocity $\vec{v}_L$:

$$\Pi'_{ij} = P \delta_{ij} + \rho (v_i - v_{Li})(v_j - v_{Lj}) . \quad (7)$$

The momentum flux through the cylindrical surface of radius r_0 is given by the integral $\int dS_j \Pi'_{ij}$ where dS_j are the components of the vector $d\vec{S}$ directed along the outer normal to the boundary of the cylindrical region and equal




Figure 1: Magnus (lift) force. It is derived from the momentum balance inside a cylinder of radius r_0 .

to the elementary area of the boundary in magnitude. Then using Eqs. (1), (5), and (7), the momentum balance yields the following relation:

$$\rho[(\vec{v}_L - \vec{v}_{tr}) \times \vec{\kappa}] = \vec{F} . \quad (8)$$

On the left-hand side of this equation one can see the Magnus force as it comes in the classical hydrodynamics. The Magnus force balances the resultant external force $\vec{F}$ applied to the cylinder. In the absence of external forces the cylinder moves with the transport velocity of the liquid: $\vec{v}_L = \vec{v}_{tr}$ (Helmholtz's theorem).

In the derivation we used the hydrodynamic equations only at large distance from the vortex line, and the radius of the cylinder does not appear in the final result. Therefore Eq. (8) is valid even if the circular flow with circulation κ occurs without any cylinder at all. In hydrodynamics such a flow pattern is called a vortex tube, a vortex line, or simply a vortex. Hydrodynamics is invalid at small distance from the vortex line. This area is called the vortex core. But this does not invalidate the derivation of the Magnus force for a Galilean invariant liquid, in which the momentum is a well-defined *conserved* quantity even inside the vortex core where the hydrodynamic theory does not hold.

In the momentum balance in a cylinder around a vortex, a half of the Magnus force is due to the Bernoulli contribution to the pressure, Eq. (5); another half is due to the convection term $\propto v_i v_j$ in the momentum flux. However, such decomposition of the resultant momentum flux onto the Bernoulli and the convection parts is not universal and depends on the shape of the area for which we consider the momentum balance. We may consider the momentum balance in a stripe, which contains the vortex inside and is oriented normally to the transport flow (see Fig. 2). This yields again Eq. (8), but now the pressure (the Bernoulli term) does not contribute to the transverse force at all, and only convection is responsible for the Magnus force. Then the physical origin of the Magnus force is the following. The liquid enters the stripe, which contains a vortex, with one value of the transverse velocity (equal to zero in Fig. 2) and exits from the stripe with another value of it. A difference between two values is $\Delta\vec{v}$ in Fig. 2. The transverse force is the total variation of the transverse (with respect to the incident velocity $\vec{v}$) liquid momentum per unit time. The latter is equal to a product of the current circulation $\rho \oint \Delta\vec{v} \cdot d\vec{l}$ and the velocity v .




Figure 2: Momentum balance in a stripe area. The transverse force is determined by the current circulation $\rho \oint \Delta \vec{v} \cdot d\vec{l}$ round the stripe.

3 The Magnus force in a superfluid

In the superfluid state liquid motion is described by two-fluid hydrodynamics: the liquid consists of the superfluid and the normal component with the superfluid and the normal density ρ_s and ρ_n , and the superfluid and the normal velocity $\vec{v}_s$ and $\vec{v}_n$ respectively. The circular motion around the vortex line is related to the superfluid motion. Therefore Hall and Vinen [2] suggested that the Magnus force is entirely connected with the superfluid density ρ_s and the superfluid velocity $\vec{v}_s(\vec{r}) = \vec{v}_v(\vec{r}) + \vec{v}_{\text{str}}$. Then instead of Eq. (8) one has:

$$\rho_s[(\vec{v}_L - \vec{v}_s) \times \vec{\kappa}] = \vec{F}. \quad (9)$$

Here and later on we omit the subscript “tr” replacing $\vec{v}_{\text{str}}$ by $\vec{v}_s$. But one should remember that the superfluid velocity $\vec{v}_s$ in the expression for the Magnus force is the superfluid velocity far from the vortex line.

The “external” force $\vec{F}$ is in fact external not for the whole liquid, but only for its superfluid part. The force appears due to interaction with quasi-particles which constitute the normal part of the liquid, and therefore is proportional to the relative velocity $\vec{v}_L - \vec{v}_n$ [2]. For a Galilean invariant

liquid with axial symmetry the most general expression for the force $\vec{F}$ is

$$\vec{F} = -D(\vec{v}_L - \vec{v}_n) - D'[\hat{z} \times (\vec{v}_L - \vec{v}_n)] . \quad (10)$$

The force component $\propto D'$ transverse to the velocity $\vec{v}_L - \vec{v}_n$ is possible because of broken time invariance in presence of a vortex and resulting asymmetry of quasiparticle scattering by a vortex.

Inserting the force $\vec{F}$ into Eq. (9), one can rewrite the equation of vortex motion collecting together the terms proportional to the velocity $\vec{v}_L$:

$$\rho_M[\vec{v}_L \times \vec{\kappa}] + D\vec{v}_L = \rho_s[\vec{v}_s \times \vec{\kappa}] + D\vec{v}_n + D'[\hat{z} \times \vec{v}_n] . \quad (11)$$

The forces on the left-hand side of the equation are the *effective Magnus force* $\propto \rho_M = \rho_s - D'/\kappa$ and the longitudinal friction force $\propto D$. The forces on the right-hand side are driving forces produced by the superfluid and normal flows. In the theory of superconductivity the force $\rho_s[\vec{v}_s \times \vec{\kappa}]$, proportional to the superfluid velocity $\vec{v}_s$, is called the Lorentz force. The left-hand side of Eq. (11) presents the response of the vortex to these driving forces. The factor ρ_M , which determines the amplitude of the effective Magnus force on the vortex, is not equal to the superfluid density ρ_s in general. In the next sections we shall consider the contribution to D' from phonon scattering by a vortex (the Iordanskii force). The contribution to D' from the bound states in the vortex core is discussed by Kopnin [14].

4 Nonlinear Schrödinger equation and two-fluid hydrodynamics

In the Gross–Pitaevskii theory [23] the ground state and weakly excited states of a Bose gas are described by the condensate wave function $\psi = a \exp(i\phi)$ which is a solution of the nonlinear Schrödinger equation

$$i\hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \psi + V|\psi|^2 \psi . \quad (12)$$

Here V is the amplitude of two-particle interaction. The nonlinear Schrödinger equation is the Euler–Lagrange equation for the Lagrangian

$$L = \frac{i\hbar}{2} \left(\psi^* \frac{\partial \psi}{\partial t} - \psi \frac{\partial \psi^*}{\partial t} \right) - \frac{\hbar^2}{2m} |\nabla \psi|^2 - \frac{V}{2} |\psi|^4 . \quad (13)$$

Noether's theorem yields the momentum conservation law

$$\frac{\partial j_i}{\partial t} + \nabla_j \Pi_{ij} \quad (14)$$

where

$$\vec{j} = -\frac{\partial L}{\partial \dot{\psi}} \vec{\nabla} \psi - \frac{\partial L}{\partial \dot{\psi}^*} \dot{\psi}^* = -\frac{i\hbar}{2} (\psi^* \vec{\nabla} \psi - \psi \vec{\nabla} \psi^*) \quad (15)$$

is the mass current and

$$\begin{aligned} \Pi_{ij} &= -\frac{\partial L}{\partial \nabla_j \psi} \nabla_i \psi - \frac{\partial L}{\partial \nabla_j \psi^*} \nabla_i \psi^* + L \delta_{ij} \\ &= \frac{\hbar^2}{2m} (\nabla_i \psi \nabla_j \psi^* + \nabla_i \psi^* \nabla_j \psi) + \delta_{ij} P \end{aligned} \quad (16)$$

is the momentum-flux tensor. The pressure P in this expression corresponds to the general thermodynamic definition of the pressure via a functional derivation of the energy with respect to the particle density $n = |\psi|^2$:

$$P = L = n \frac{\delta E}{\delta n} - E = n \left[\frac{\partial E}{\partial n} - \vec{\nabla} \cdot \left(\frac{\partial E}{\partial \vec{\nabla} n} \right) \right] - E = \frac{V}{2} |\psi|^4 - \frac{\hbar^2}{4m} \nabla^2 |\psi|^2, \quad (17)$$

where

$$E = \frac{\partial L}{\partial \dot{\psi}} \dot{\psi} + \frac{\partial L}{\partial \dot{\psi}^*} \dot{\psi}^* - L = \frac{\hbar^2}{2m} |\nabla \psi|^2 + \frac{V}{2} |\psi|^4 \quad (18)$$

is the energy density. But in the hydrodynamic limit (see below) the dependence of the energy on the density gradient is usually neglected.

Using the Madelung transformation [28], the nonlinear Schrödinger equation (12) for a complex function may be transformed into two real equations for the liquid density $\rho = ma^2$ and the liquid velocity $\vec{v} = (\kappa/2\pi) \vec{\nabla} \phi$ where $\kappa = h/m$ is the circulation quantum. Far from the vortex line these equations are hydrodynamic equations for an ideal inviscous liquid:

$$\frac{\partial \rho}{\partial t} + \vec{\nabla} \cdot (\rho \vec{v}) = 0, \quad (19)$$

$$\frac{\partial \vec{v}}{\partial t} + (\vec{v} \cdot \vec{\nabla}) \vec{v} = -\vec{\nabla} \mu. \quad (20)$$

Here $\mu = Va^2/m$ is the chemical potential. Equation (16) becomes the hydrodynamic momentum-flux tensor $\Pi_{ij} = P\delta_{ij} + \rho v_i(\vec{r})v_j(\vec{r})$. Thus the hydrodynamics of an ideal liquid directly follows from the nonlinear Schrödinger equation.

Suppose that a plane sound wave propagates in the liquid generating the phase variation $\phi(\vec{r}, t) = \phi_0 \exp(i\vec{k} \cdot \vec{r} - i\omega t)$. Then the liquid density and velocity are functions of the time t and the position vector $\vec{r}$ in the plane xy :

$$\rho(\vec{r}, t) = \rho_0 + \rho_{(1)}(\vec{r}, t) , \quad \vec{v}(\vec{r}, t) = \vec{v}_0 + \vec{v}_{(1)}(\vec{r}, t) , \quad (21)$$

where ρ_0 and $\vec{v}_0$ are the average density and the average velocity in the liquid, $\rho_{(1)}(\vec{r}, t)$ and $\vec{v}_{(1)}(\vec{r}, t) = \frac{\kappa}{2\pi} \vec{\nabla} \phi$ are periodical variations of the density and the velocity due to the sound wave ($\langle \rho_{(1)} \rangle = 0$, $\langle \vec{v}_{(1)} \rangle = 0$). They should be determined from Eqs. (19) and (20) after their linearization. In particular, Eq. (20) gives the relation between the density variation and the phase ϕ :

$$\rho_{(1)} = \frac{\rho_0}{c_s^2} \mu_{(1)} = -\frac{\rho_0}{c_s^2} \frac{\kappa}{2\pi} \left\{ \frac{\partial \phi}{\partial t} + \vec{v}_0 \cdot \vec{\nabla} \phi(\vec{r}) \right\} , \quad (22)$$

where $c_s = \sqrt{Va^2/m}$ is the sound velocity. Substitution of this expression into Eq. (19) yields the sound equation for a moving liquid with the wave spectrum $\omega = c_s k + \vec{k} \cdot \vec{v}_0$.

The sound propagation is accompanied with the transport of mass. This is an effect of the second order with respect to the wave amplitude. The total mass current expanded with respect to the wave amplitude and averaged over time is

$$\vec{j} = \rho_0 \vec{v}_0 + \langle \rho_{(1)} \vec{v}_{(1)} \rangle + \langle \rho_{(2)} \rangle \vec{v}_0 + \rho_0 \langle \vec{v}_{(2)} \rangle , \quad (23)$$

where

$$\langle \rho_{(1)} \vec{v}_{(1)} \rangle = \vec{j}^{ph}(\vec{p}) = \rho_0 \phi_0^2 \frac{\kappa^2 k}{8\pi^2 c_s} \vec{k} = n(\vec{p}) \vec{p} . \quad (24)$$

is the mass current and $n(\vec{p})$ is the number of phonons with the momentum $\vec{p} = \hbar \vec{k}$ and the energy $E = \varepsilon(\vec{p}) + \vec{p} \cdot \vec{v}_0$. The phonon mass current is the phonon momentum density in the reference frame moving with the average liquid velocity $\vec{v}_0$, and $\varepsilon(\vec{p}) = c_s p$ is the energy in the same reference frame.

Mathematically the second-order corrections to the mass density, $\langle \rho_{(2)} \rangle$, and the average velocity, $\langle \vec{v}_{(2)} \rangle$, remain undefined, but there are physical constraints to specify them. First of all, we assume that phonon excitations

do not change the average mass density, and $\langle \rho_{(2)} \rangle$ must vanish. As for the second-order correction $\langle \vec{v}_{(2)} \rangle$ to the average velocity, it should produce a second-order correction $\langle \phi_{(2)} \rangle$ to the phase which is impossible in quantum hydrodynamics. The simplest way to see it is to consider the propagation of phonons in an annular channel with the periodic boundary conditions. The phase variation over the channel length is a topological invariant, and weak excitations (phonons) cannot change it. Therefore $\langle \vec{v}_{(2)} \rangle$ must vanish. More complicated arguments must be given for an open geometry, but intuitively it is clear that this basic physical constrain should not depend on the boundary condition.

It is important to emphasize a difference between sound waves in a liquid and in an elastic solid. The sound wave in the elastic solid is not accompanied by real mass transport: all atoms oscillate near their equilibrium positions in the crystal lattice, but they cannot move in average if the crystal is fixed at a laboratory table. Within our present formalism this means that the second-order contribution $\langle \vec{v}_{(2)} \rangle$ to the average velocity must not vanish, but compensate the second-order contribution $\langle \rho_{(1)} \vec{v}_{(1)} \rangle$ to the mass current. Finally a phonon in a solid cannot have a real momentum but only a quasimomentum. The problem of the phonon momentum in liquids and solids has already been discussed a long time [29] (see also the recent paper by Stone [30]), and they have noticed that the sound wave may have a different momentum using Euler or Lagrange variables. In fact, the momentum should not depend on a choice of variables, but only on physical conditions. However, at various physical conditions a proper choice of variables can do an analysis more straightforward. In solids the Lagrange variables are preferable since in this case the velocity is related to a given particle and coincides with the *center-of-mass* velocity which must not change in average and therefore has no second-order corrections. Using Euler variables in liquids the *average* velocity $\vec{v}_0$ relates to a given point in the space and has no second-order corrections due to phonons.

Expanding the momentum-flux tensor up to the terms of the second order with respect to the sound wave amplitude one obtains:

$$\Pi_{ij} = P_0 \delta_{ij} + \rho_0 v_{0i} v_{0j} + \Pi_{ij}^{ph} , \quad (25)$$

where the second-order phonon contribution is

$$\Pi_{ij}^{ph} = \langle P_{(2)} \rangle \delta_{ij} + \langle \rho_{(1)} (v_{(1)})_i \rangle v_{0j} + \langle \rho_{(1)} (v_{(1)})_j \rangle v_{0i} + \rho_0 \langle (v_{(1)})_i (v_{(1)})_j \rangle . \quad (26)$$

The second-order contribution $P_{(2)}$ to the pressure can be obtained from the Gibbs-Duhem relation $\delta P = \rho \delta \mu$ at $T = 0$ using expansions $\rho = \rho_0 + \rho_{(1)}$ and $\mu = \mu_0 + \mu_{(1)} + \mu_{(2)}$, where μ_0 is the chemical potential without the sound wave. This yields $P_{(2)} = \rho_0 \mu_{(2)} + \frac{\partial \rho}{\partial \mu} \frac{\rho_{(1)}^2}{2}$, where $\frac{\partial \rho}{\partial \mu} = \rho_0 / c_s^2$. According to the Euler equation (3) the second-order contribution to the chemical potential is $\mu_{(2)} = -\frac{v_{(1)}^2}{2}$. Then

$$\langle P_{(2)} \rangle = \frac{c_s^2}{\rho_0} \frac{\langle \rho_{(1)}^2 \rangle}{2} - \rho_0 \frac{\langle v_{(1)}^2 \rangle}{2} . \quad (27)$$

In the presence of the phonon distribution the total mass current is

$$\vec{j} = \rho_0 \vec{v}_0 + \frac{1}{h^3} \int \vec{j}^{ph}(\vec{p}) d_3 \vec{p} = \rho_0 \vec{v}_0 + \frac{1}{h^3} \int n(\vec{p}) \vec{p} d_3 \vec{p} , \quad (28)$$

In the thermal equilibrium at $T > 0$, the phonon numbers are given by the Planck distribution $n(\vec{p}) = n_0(E, \vec{v}_n)$ with the drift velocity $\vec{v}_n$ of quasiparticles:

$$n_0(E, \vec{v}_n) = \frac{1}{\exp \frac{E(\vec{p}) - \vec{p} \cdot \vec{v}_n}{T} - 1} = \frac{1}{\exp \frac{\varepsilon(\vec{p}) + \vec{p} \cdot (\vec{v}_0 - \vec{v}_n)}{T} - 1} . \quad (29)$$

Linearizing Eq. (29) with respect to the relative velocity $\vec{v}_0 - \vec{v}_n$, one obtains from Eq. (28) that

$$\vec{j} = \rho_0 \vec{v}_0 + \rho_n (\vec{v}_n - \vec{v}_0) . \quad (30)$$

This expression is equivalent to the two-fluid expression $\vec{j} = \rho \vec{v}_s + \rho_n (\vec{v}_n - \vec{v}_s) = \rho_s \vec{v}_s + \rho_n \vec{v}_n$ assuming that $\rho = \rho_0 = \rho_s + \rho_n$, $\vec{v}_0 = \vec{v}_s$, and the normal density is given by the usual two-fluid-hydrodynamics expression:

$$\rho_n = -\frac{1}{3h^3} \int \frac{\partial n_0(\varepsilon, 0)}{\partial E} p^2 d_3 \vec{p} . \quad (31)$$

In the same manner one can derive the two-fluid momentum-flux tensor [7]:

$$\Pi_{ij} = P \delta_{ij} + \rho_s v_{si} v_{sj} + \rho_n v_{ni} v_{nj} . \quad (32)$$

This analysis demonstrates that two-fluid-hydrodynamics relations can be derived from the hydrodynamics of an ideal liquid in presence of thermally excited sound waves, as was shown by Putterman and Roberts [31]. In order to obtain a complete system of equations of the two-fluid theory, one should take into account phonon-phonon interaction, which is essential for

the phonon distribution function being close to the equilibrium Planck distribution. In the two-fluid theory the locally defined superfluid and normal velocities $\vec{v}_s$ and $\vec{v}_n$ correspond to the average velocity of a liquid in a fixed point of the space and to the drift velocity of the phonon gas respectively. The two-fluid hydrodynamics is valid only at scales exceeding the phonon mean-free path l_{ph} .

5 Scattering of phonons by the vortex in hydrodynamics

Phonon scattering by a vortex line was studied beginning from the works by Pitaevskii [32] and Fetter [33]. We consider a sound wave propagating in the plane xy normal to a vortex line (the axis z). In the linearized hydrodynamic equations of the previous section the fluid velocity $\vec{v}_0$ should be replaced by the velocity $\vec{v}_v(\vec{r})$ around the vortex line:

$$\frac{\partial \rho_{(1)}}{\partial t} + \rho_0 \vec{\nabla} \cdot \vec{v}_{(1)} = -\vec{v}_v \cdot \vec{\nabla} \rho_{(1)} , \quad (33)$$

$$\frac{\partial \vec{v}_{(1)}}{\partial t} + \frac{c_s^2}{\rho_0} \vec{\nabla} \rho_{(1)} = - \left[(\vec{v}_v \cdot \vec{\nabla}) \vec{v}_{(1)} + (\vec{v}_{(1)} \cdot \vec{\nabla}) \vec{v}_v \right] . \quad (34)$$

Using the vector identity

$$(\vec{v} \cdot \vec{\nabla}) \vec{v} = \vec{\nabla} \frac{v^2}{2} - \vec{v} \times [\vec{\nabla} \times \vec{v}] \quad (35)$$

for the velocity $\vec{v} = \vec{v}_v + \vec{v}_{(1)}$, Eq. (34) can be rewritten as

$$\frac{\partial \vec{v}_{(1)}}{\partial t} + \frac{c_s^2}{\rho_0} \vec{\nabla} \rho_{(1)} = -\vec{\nabla} (\vec{v}_v \cdot \vec{v}_{(1)}) + [\vec{v}_{(1)} \times \vec{\kappa}] \delta_2(\vec{r}) . \quad (36)$$

The perturbation from the vortex (the right-hand side) contains a δ -function because the vortex line is not at rest when the sound wave propagates past the vortex. In order to weaken the singularity one can introduce the time-dependent vortex velocity $\vec{v}_v(\vec{r}, t)$ as a zero-order approximation for the velocity field [5]. Then $\vec{r}$ in Eq. (1) must be replaced by $\vec{r} - \vec{v}_L t$ and $\partial \vec{v}_v / \partial t = -(\vec{v}_L \cdot \vec{\nabla}) \vec{v}_v = -\vec{\nabla} (\vec{v}_L \cdot \vec{v}_v) + [\vec{v}_L \times \vec{\kappa}] \delta_2(\vec{r})$. Since there is no external

force on the liquid, the vortex moves with the velocity in the sound wave: $\vec{v}_L = \vec{v}_{(1)}(0, t)$. Now in the linearization procedure the fluid acceleration in Eq. (20) must be presented as $\partial\vec{v}/\partial t = \partial\vec{v}_v/\partial t + \partial\vec{v}_{(1)}/\partial t$. As a result Eq. (36) is replaced by:

$$\frac{\partial\vec{v}_{(1)}}{\partial t} + \frac{c_s^2}{\rho_0} \vec{\nabla} \rho_{(1)} = \vec{\nabla}[\vec{v}_v \cdot \vec{v}_{(1)}(\vec{r})] - \vec{\nabla}[\vec{v}_v \cdot \vec{v}_{(1)}(0)] . \quad (37)$$

Equation (37) yields:

$$\rho_{(1)} = -\frac{\rho_0}{c_s^2} \frac{\kappa}{2\pi} \left\{ \frac{\partial\phi}{\partial t} + \vec{v}_v \cdot [\vec{\nabla}\phi(\vec{r}) - \vec{\nabla}\phi(0)] \right\} . \quad (38)$$

Substitution of $\rho_{(1)}$ in Eq. (33) yields the linear sound equation for the phase:

$$\frac{\partial^2\phi}{\partial t^2} - c_s^2 \vec{\nabla}^2 \phi = -2\vec{v}_v(\vec{r}) \cdot \vec{\nabla} \frac{\partial}{\partial t} \left[\phi(\vec{r}) - \frac{1}{2}\phi(0) \right] . \quad (39)$$

In the long-wavelength limit $k \rightarrow 0$ one may use the Born approximation. The Born perturbation parameter $\kappa k/c_s$ is on the order of the ratio of the vortex core radius $r_c \sim \kappa/c_s$ to the wavelength $2\pi/k$. Then after substituting the plane wave into the right-hand side of Eq. (39) the solution of this equation is

$$\phi = \phi_0 \exp(-i\omega t) \left\{ \exp(i\vec{k} \cdot \vec{r}) - \frac{ik}{4c_s} \int d_2\vec{r}_1 H_0^{(1)}(k|\vec{r} - \vec{r}_1|) \vec{k} \cdot \vec{v}_v(\vec{r}_1) [2 \exp(i\vec{k} \cdot \vec{r}_1) - 1] \right\} . \quad (40)$$

Here $H_0^{(1)}(z)$ is the zero-order Hankel function of the first kind, and $\frac{i}{4}H_0^{(1)}(k|\vec{r} - \vec{r}_1|)$ is the Green function for the 2D wave equation, i. e., satisfies to the equation

$$-(k^2 + \vec{\nabla}^2)\phi(\vec{r}) = \delta_2(\vec{r} - \vec{r}_1) . \quad (41)$$

In the standard scattering theory they use the asymptotic expression for the Hankel function at large values of the argument:

$$\lim_{z \rightarrow \infty} H_0^{(1)}(z) = \sqrt{\frac{2}{\pi z}} e^{i(z - \pi/4)} . \quad (42)$$

If the perturbation is confined to a vicinity of the line, then $r_1 \ll r$ and

$$|\vec{r} - \vec{r}_1| \approx r - \frac{(\vec{r}_1 \cdot \vec{r})}{r} . \quad (43)$$

After integration in Eq. (40) the wave at $kr \gg 1$ becomes a superposition of the incident plane wave $\propto \exp(i\vec{k} \cdot \vec{r})$ and the scattered wave $\propto \exp(ikr)$:

$$\phi = \phi_0 \exp(-i\omega t) \left[\exp(i\vec{k} \cdot \vec{r}) + \frac{ia(\varphi)}{\sqrt{r}} \exp(ikr) \right] . \quad (44)$$

Here $a(\varphi)$ is the scattering amplitude which is a function of the angle φ between the initial wave vector $\vec{k}$ and the wave vector $\vec{k}' = k\vec{r}/r$ after scattering (see Fig. 3). In the Born approximation (see the paper [7] and references therein for more details)

$$a(\varphi) = \sqrt{\frac{k}{2\pi}} \frac{1}{c_s} e^{i\frac{\pi}{4}} [\hat{k} \times \vec{k}'] \cdot \vec{k} \frac{1}{q^2} \left(1 - \frac{q^2}{2k^2} \right) = -\frac{1}{2} \sqrt{\frac{k}{2\pi}} \frac{\kappa}{c_s} e^{i\frac{\pi}{4}} \frac{\sin \varphi \cos \varphi}{1 - \cos \varphi} , \quad (45)$$

where $\vec{q} = \vec{k} - \vec{k}'$ is the momentum transferred by the scattered phonon to the vortex, and $q^2 = 2k^2(1 - \cos \varphi)$.

Thus the vortex is a line defect which scatters a sound wave. Scattering produces a force on the defect (vortex). If the perturbation by the line defect is confined to a finite vicinity of the line, the force

$$\vec{F}^{ph} = \sigma_{\parallel} c_s \vec{j}^{ph} - \sigma_{\perp} c_s [\hat{z} \times \vec{j}^{ph}] \quad (46)$$

is determined by two effective cross-sections [5, 7]: the transport cross-section for the dissipative force,

$$\sigma_{\parallel} = \int \sigma(\varphi) (1 - \cos \varphi) d\varphi , \quad (47)$$

and the transverse cross-section for the transverse force,

$$\sigma_{\perp} = \int \sigma(\varphi) \sin \varphi d\varphi . \quad (48)$$

The differential cross-section $\sigma(\varphi) = |a(\varphi)|^2$ with $a(\varphi)$ from Eq. (45) is quadratic in the circulation κ and even in φ . Therefore in the Born approximation the transverse cross-section $\sigma_{\perp}$ vanishes.




Figure 3: Scattering of a sound wave by a vortex.

However, due to a very slow decrease of the velocity $v_v \propto 1/r$ far from the vortex, the scattering amplitude is divergent at small scattering angles $\varphi \rightarrow 0$:

$$\lim_{\varphi \rightarrow 0} a(\varphi) = -\sqrt{\frac{k}{2\pi}} \frac{\kappa}{c_s} e^{i\frac{\pi}{4}} \frac{1}{\varphi}. \quad (49)$$

This divergence is integrable in the integral for the transport cross-section, Eq. (47), and its calculation is reliable. Contrary to it, the integrand in Eq. (48) for the transverse cross-section has a pole at $\varphi = 0$. The principal value of the integral vanishes, but there is no justification for the choice of the principal value, and the contribution of the small angles requires an additional analysis.

At small scattering angles $\varphi \ll 1/\sqrt{kr}$ the asymptotic expansion Eq. (44) is invalid, and one cannot use the differential cross-section or the scattering amplitude for description of the small-angle scattering [4, 5, 9, 34]. Meanwhile, the small-angle behavior is crucial for the transverse force as demonstrated below. The accurate calculation of the integral in Eq. (40) for

small angles (see Ref. [5] and Appendix B in Ref. [7]) yields that at $\varphi \ll 1$

$$\phi = \phi_0 \exp(-i\omega t + i\vec{k} \cdot \vec{r}) \left[1 + \frac{i\kappa k}{2c_s} \Phi \left(\varphi \sqrt{\frac{kr}{2i}} \right) \right] . \quad (50)$$

Using an asymptotic expression for the error integral

$$\Phi(z) = \frac{2}{\sqrt{\pi}} \int_0^z e^{-t^2} dt \longrightarrow \frac{z}{|z|} - \frac{1}{\sqrt{\pi}z} \exp(-z^2) \quad (51)$$

at $|z| \rightarrow \infty$, one obtains for angles $1 \gg \varphi \gg 1/\sqrt{kr}$:

$$\phi = \phi_0 \exp(-i\omega t) \left[\exp(i\vec{k} \cdot \vec{r}) \left(1 + \frac{i\kappa k}{2c_s} \frac{\varphi}{|\varphi|} \right) - \frac{i\kappa}{c_s} \sqrt{\frac{k}{2\pi r}} \frac{1}{\varphi} \exp \left(ikr + i\frac{\pi}{4} \right) \right] . \quad (52)$$

The second term in square brackets coincides with scattering wave at small angles $\varphi \ll 1$ with the amplitude given by Eq. (49). But now one can see that the standard scattering theory misses to reveal an important non-analytical correction to the incident plane wave, which changes a sign when the scattering angle φ crosses zero. Its physical meaning is discussed in the next section.

6 The Iordanskii force and the Aharonov–Bohm effect

The analogy between the phonon scattering by a vortex and the Aharonov–Bohm effect for electrons scattered by a magnetic-flux tube becomes evident if one rewrites the sound equation (39) in presence of the vortex as

$$k^2 \phi - \left(-i\vec{\nabla} + \frac{k}{c_s} \vec{v}_v \right)^2 \phi = 0 . \quad (53)$$

It differs from the sound equation (39) by the term of the second order in $v_v \propto \kappa$ and by absence of the contribution from the vortex-line motion [the term $\propto \phi(0)$ on the right-hand side of Eq. (39)]. This difference is unimportant for the calculation of the transverse force, which is linear in κ .

On the other hand, the stationary Schrödinger equation for an electron in presence of the magnetic flux Φ confined to a thin tube (the Aharonov–Bohm effect [8]) is:

$$E\psi(\vec{r}) = \frac{1}{2m} \left(-i\hbar\vec{\nabla} - \frac{e}{c}\vec{A} \right)^2 \psi(\vec{r}) . \quad (54)$$

Here ψ is the electron wave function with energy E and the electromagnetic vector potential is connected with the magnetic flux Φ by the relation similar to that for the velocity $\vec{v}_v$ around the vortex line [Eq. (1)]:

$$\vec{A} = \Phi \frac{[\hat{z} \times \vec{r}]}{2\pi r^2} . \quad (55)$$

Let us consider the quasiclassical solution of the sound equation:

$$\phi = \phi_0 \exp \left(-i\omega t + i\vec{k} \cdot \vec{r} + \frac{i\delta S}{\hbar} \right) = \phi_0 \exp(-i\omega t + i\vec{k} \cdot \vec{r}) \left(1 + \frac{i\kappa k}{2\pi c_s} \theta \right) , \quad (56)$$

where $\delta S = -(\hbar k/c_s) \int^{\vec{r}} \vec{v}_v \cdot d\vec{l} = \hbar \theta \kappa k / 2\pi c_s$ is the variation of the action due to interaction with the circular velocity from the vortex along quasiclassical trajectories. The angle θ is an azimuth angle for the position vector $\vec{r}$ measured from the direction opposite to the wave vector $\vec{k}$ (see Fig. 3). This choice provides that the quasiclassical correction vanishes for the incident wave far from the vortex. One can check directly that Eq. (56) satisfies the sound equation (39) in the first order of the parameter $\kappa k/c_s$. For the Aharonov–Bohm effect the phase $\delta S/\hbar$ arises from the electromagnetic vector potential: $\delta S = -(e/c) \int^{\vec{r}} \vec{A} \cdot d\vec{l}$.

The velocity generated by the sound wave around the vortex is

$$\vec{v}_{(1)} = \frac{\kappa}{2\pi} \vec{\nabla} \phi = \frac{\kappa}{2\pi} \phi_0 \exp(-i\omega t + i\vec{k} \cdot \vec{r}) \left(i\vec{k} - \frac{ik}{c_s} \vec{v}_v \right) . \quad (57)$$

From Eqs. (38) and (57) we can obtain the phonon mass current with first-order corrections in $\vec{v}_v$.

$$\vec{j}_{ph} = \langle \rho_{(1)} \vec{v}_{(1)} \rangle = \frac{1}{8\pi^3} \int d\vec{k} n(\hbar\vec{k}) \hbar \left[\vec{k} - \frac{k}{c_s} \vec{v}_v(\vec{r}) - (\vec{k} \cdot \vec{v}_v(\vec{r})) \frac{\vec{k}}{c_s k} \right] . \quad (58)$$

Due to the last term in this expression, the phonon mass current is not curl-free.




Figure 4: Aharonov–Bohm interference.

According to Eq. (56) the phase ϕ is multivalued, and one must choose a cut for an angle θ at the direction $\vec{k}$, where $\theta = \pm\pi$. The jump of the phase on the cut line behind the vortex is a manifestation of the Aharonov–Bohm effect [8]: the sound wave after its interaction with the vortex has different phases on the left and on the right of the vortex line. This results in an interference [5, 7].

In the interference region the quasiclassical solution is invalid and must be replaced by Eq. (50). The width of the interference region is $d_{int} \sim \sqrt{r/k}$. Here r is the distance from the vortex line. The interference region corresponds to very small scattering angles $\sim d_{int}/r = 1/\sqrt{kr}$.

Now we are ready to consider the momentum balance using the condition that $\int dS_j \Pi_{\perp j} = 0$ where subscript $\perp$ is for a component normal to the wave vector $\vec{k}$ of the incident wave. The total momentum-flux tensor can be obtained from Eqs. (25) and (26) assuming $\vec{v}_0(\vec{r}) = \vec{v}_v(\vec{r}) + \vec{v}_s$ and neglecting

some unimportant terms:

$$\begin{aligned} \Pi_{ij} = & -\rho_0(\vec{v}_s - \vec{v}_L) \cdot \vec{v}_v \delta_{ij} + \rho_0 v_{0i} v_{0j} \\ & + \langle \rho_{(1)}(v_{(1)})_i \rangle v_{vj} + \langle \rho_{(1)}(v_{(1)})_j \rangle v_{vi} + \rho_0 \langle (v_{(1)})_i (v_{(1)})_j \rangle . \end{aligned} \quad (59)$$

The first two terms in this expression yield the momentum flux without phonons, which gives the Magnus force for a liquid with the density ρ_0 and the velocity $\vec{v}_s$. The term $\propto v_{vj}$ does not contribute to the momentum flux through a cylindrical surface around the vortex, since the velocity $\vec{v}_v$ is tangent to this surface. The term $v_{vi} \langle \rho_{(1)}(v_{(1)})_j \rangle$, in which the mass current $\langle \rho_{(1)}(v_{(1)})_j \rangle$ is given Eq. (24) for the plane wave in absence of the vortex, gives a contribution to the momentum flux, which exactly cancels the contribution of the term $\rho_0 \langle (v_{(1)})_i (v_{(1)})_j \rangle$ outside the interference region, where the velocity is given by Eq. (57). Finally only the interference region contributes the momentum flux $\int dS_j \Pi_{\perp j}$.

In the interference region the velocity is obtained by taking the gradient of the phase given by Eq. (50). Its component normal to $\vec{k}$,

$$v_{(1)\perp} = \frac{\kappa}{2\pi r} \frac{\partial \phi}{\partial \varphi} = \phi_0 \exp(-i\omega t + i\vec{k} \cdot \vec{r}) \frac{i\kappa^2 k}{4\pi c_s r} \frac{\partial \Phi(\varphi \sqrt{kr/2i})}{\partial \varphi} , \quad (60)$$

determines the interference contribution to the transverse force:

$$\begin{aligned} \int dS_j \rho_0 \langle (v_{(1)})_{\perp} (v_{(1)})_j \rangle &= \int \rho_0 \langle (v_{(1)})_{\perp} (v_{(1)})_r \rangle r d\varphi \\ &= \frac{1}{8\pi^2} \rho_0 \phi_0^2 \frac{\kappa^2 k}{\hbar} [\delta S_- - \delta S_+] = \frac{1}{8\pi^2} \rho_0 \phi_0^2 \frac{\kappa^3 k^2}{c_s} = \kappa j^{ph} , \end{aligned} \quad (61)$$

where $\delta S_{\pm}$ are the action variations at $\theta \rightarrow \mp\pi$. Thus the interference region, which corresponds to an infinitesimally small angle interval, yields a finite contribution to the transverse force, which one could not obtain from the standard scattering theory using the differential cross-section. In fact the details of the solution in the interference region are not essential: only a jump of the phase across the interference region is of importance.

Finally the momentum balance condition $\int dS_j \Pi_{\perp j} = 0$ yields the relation

$$\rho_0[(\vec{v}_L - \vec{v}_s) \times \vec{\kappa}] - [\vec{j}^{ph}(\vec{p}) \times \vec{\kappa}] = 0 . \quad (62)$$

The second vector product on the left-hand side is a transverse force which corresponds to the transverse cross-section [see Eq. (48)]

$$\sigma_{\perp} = \frac{\delta S_{-} - \delta S_{+}}{\hbar k} \quad (63)$$

equal to κ/c_s . For the Planck distribution of phonons, $\vec{j}^{ph}$ must be replaced by $\rho_n(\vec{v}_n - \vec{v}_s)$:

$$\rho_s[(\vec{v}_L - \vec{v}_s) \times \vec{\kappa}] + \rho_n[(\vec{v}_L - \vec{v}_n) \times \vec{\kappa}] = 0. \quad (64)$$

The term $\propto (\vec{v}_L - \vec{v}_n)$ is the Iordanskii force, which corresponds to $D' = -\kappa\rho_n$ in Eq. (10). The longitudinal force $\propto D$ is not present in Eq. (64) since we ignored terms of the second order in κ in order to simplify our derivation. According to Eq. (64) the vortex moves with the center-of-mass velocity $\vec{v} = \frac{\rho_s}{\rho}\vec{v}_s + \frac{\rho_n}{\rho}\vec{v}_n$.

Our scattering analysis was done in the coordinate frame where the vortex is at rest and we neglected the relative superfluid velocity $\vec{v}_s - \vec{v}_L$ with respect to the vortex. If the velocity $\vec{v}_s - \vec{v}_L$ is high it can affect the value of the effective cross-section. But since the phonon momentum is linear with respect to $\vec{v}_n - \vec{v}_s$, the dependence of the cross-section on $\vec{v}_s - \vec{v}_L$ is a nonlinear effect. Thus our momentum balance took into account all effects linear in the superfluid velocity $\vec{v}_s$. This is confirmed by a more elaborate analysis of Stone [35].

7 Partial-wave analysis and the Aharonov–Bohm effect

Studying interaction of phonons with a vortex we solved the sound equation using the perturbation theory. It is completely justified because the parameter of the perturbation theory $\kappa k/c_s$ is the ratio between the vortex core and the phonon wavelength, which is always small for phonons. But in the Aharonov–Bohm problem for electrons the corresponding parameter is $\gamma = \Phi/\Phi_1$, where $\Phi_1 = hc/e$ is the magnetic-flux quantum for one electron (two times larger than the magnetic-flux quantum $\Phi_0 = hc/2e$ for a Cooper pair). This parameter can be arbitrary large, and the perturbation theory is not enough to describe an expected periodic dependence on γ . On the other

hand, there is an exact solution of the Aharonov–Bohm problem for electrons obtained by the partial-wave expansion [8], and it will be shown now how to derive the transverse force from this solution. Another derivation of the force on the Aharonov–Bohm flux tube using the wave-packet presentation is discussed by Shelankov [12].

We look for a solution of Eq. (54) as a superposition of the partial cylindrical waves $\psi = \sum_l \psi_l(r) \exp(il\varphi)$. Partial-wave amplitudes ψ_l should satisfy equations in the cylindrical system of coordinates (r, φ) :

$$\frac{d^2\psi_l}{dr^2} + \frac{1}{r} \frac{d\psi_l}{dr} - \frac{(l - \gamma)^2}{r^2} \psi_l + k^2 \psi_l = 0 . \quad (65)$$

Here k is the wave number of the electron far from the vortex so that $E = \hbar^2 k^2 / 2m$. We need a solution of Eq. (65), which at large distances has an asymptotic behavior given by Eq. (44):

$$\psi_l = \sqrt{\frac{2n}{\pi k r}} \exp \left[i \frac{\pi}{2} l + i \delta_l \right] \cos \left(k r - \frac{\pi}{2} l + \delta_l - \frac{\pi}{4} \right) , \quad (66)$$

where n is the particle density and the partial-wave phase shifts are

$$\delta_l = (l - |l - \gamma|) \pi / 2 . \quad (67)$$

For $\delta_l = 0$ Eq. (66) yields the partial-wave amplitudes of the incident plane wave at large distances r . But for nonzero δ_l there is also the scattered wave in Eq. (44) with the scattering amplitude

$$a(\varphi) = \sqrt{\frac{1}{2\pi k}} \exp \left(i \frac{\pi}{4} \right) \sum_l [1 - \exp(2i\delta_l)] \exp(il\varphi) . \quad (68)$$

The transverse force is determined by the transverse cross-section

$$\begin{aligned} \sigma_{\perp} &= \int |a(\varphi)|^2 \sin \varphi d\varphi \\ &= \frac{1}{2ik} \left\{ \sum_l e^{i2\delta_{l-1}} - \sum_l e^{i2\delta_{l+1}} + \sum_l e^{i2(\delta_{l+1}-\delta_l)} - \sum_l e^{i2(\delta_{l-1}-\delta_l)} \right\} . \end{aligned} \quad (69)$$

Shifting the number l by 2 in the first sum and by one in the fourth sum one obtains the expression for the transverse cross-section in the partial-wave

method derived long ago by Cleary [36]:

$$\sigma_{\perp} = \int |a(\varphi)|^2 \sin \varphi d\varphi = \frac{1}{k} \sum_l \sin 2(\delta_l - \delta_{l+1}) . \quad (70)$$

Using the phase shift values for the Aharonov–Bohm solution, Eq. (67), the transverse cross-section is

$$\sigma_{\perp} = -\frac{1}{k} \sin 2\pi\gamma . \quad (71)$$

However, the shift of l in the first sum of Eq. (69) is not an innocent operation because of divergence of the first and second sum at $l \rightarrow \pm\infty$. The derivation of Cleary's formula (70) assumes that the first and the second divergent sums in Eq. (69) should exactly cancel after the shift of l . But if one does not shift l , the difference of the first and the second sum is finite. Moreover, this difference cancels the contribution of the third and the fourth sum, and $\sigma_{\perp}$ vanishes in the first order with respect to the phase shifts δ_l .

Ambiguity in the calculation of the partial-wave sum for the transverse cross-section is another manifestation of the small-scattering-angle problem in the configurational space: the standard scattering theory does not provide a recipe how to treat a singularity at zero scattering angle. The zero-angle singularity is responsible for a divergent partial-wave series. The way to avoid the ambiguity is similar to that in the configurational space: one should not use the concept of the scattering amplitude for calculation of the transverse force.

We must analyze the momentum balance. The momentum-flux tensor for the electron Schrödinger equation (54) is

$$\begin{aligned} \Pi_{ij} = \frac{1}{2m} \text{Re} \left\{ \left(-i\hbar \nabla_i - \frac{e}{c} A_i \right) \psi \left(i\hbar \nabla_i - \frac{e}{c} A_i \right) \psi^* \right. \\ \left. - \psi^* \left(-i\hbar \nabla_i - \frac{e}{c} A_i \right) \left(-i\hbar \nabla_j - \frac{e}{c} A_j \right) \nabla_j \psi \right\} . \end{aligned} \quad (72)$$

If the axis x is directed along the wave vector $\vec{k}$ of the incident wave, the transverse force is determined by the momentum-flux tensor component Π_{yr} , where r is the radial coordinate in the cylindrical coordinate system $r = \sqrt{x^2 + y^2}$ and $\varphi = \arctan(y/x)$. In Eq. (72) the terms A and $\partial/r\partial\varphi$,

which are inversely proportional to r , are not important at large r . Then

$$\Pi_{yr} = \frac{\hbar^2}{2m} \sum_{l'} \sum_l \operatorname{Re} \left\{ \sin \varphi \left[\frac{\partial \psi_{l'}^*}{\partial r} \frac{\partial \psi_l}{\partial r} - \psi_l^* \frac{\partial^2 \psi_{l'}^*}{\partial r^2} \right] e^{i(l-l')\varphi} \right\} , \quad (73)$$

where ψ_l are given by Eq. (66). Finally the transverse force is

$$F_{\perp} = \oint \Pi_{yr} r d\varphi = \pi r \frac{\hbar^2}{m} \operatorname{Im} \sum_l \left\{ \frac{\partial \psi_{l+1}^*}{\partial r} \frac{\partial \psi_l}{\partial r} - \psi_l^* \frac{\partial^2 \psi_{l+1}^*}{\partial r^2} \right\} . \quad (74)$$

Here we also made a shift of l in some sums, but it is not dangerous because the terms of these sums decrease in the limits $l \rightarrow \pm\infty$.

Inserting the partial-wave amplitudes and their radial derivatives into the expression for the force we obtain:

$$F_{\perp} = -kn \frac{\hbar^2}{m} \sum_l \sin 2(\delta_{l+1} - \delta_l) = -jv\sigma_{\perp} , \quad (75)$$

where $j = \hbar kn$ and $v = \hbar k/m$ are the momentum density and the velocity in the incident plane wave, and $\sigma_{\perp}$ is the effective cross-section Eq. (70) derived by Cleary [36]. Thus there is a well-defined effective transverse cross-section, despite that we cannot use directly its expression (48) via the differential cross-section. But formally we can “repair” this expression by a recipe how to treat the singularity at small angles: we should add to the differential cross-section a singular term proportional to the derivative of $\delta(\varphi)$, and to take the principle value of the integral over the rest part of the differential cross-section.

One can obtain the transverse cross-section for phonons directly from Eq. (70) assuming that $\sin 2\pi\gamma \approx 2\pi\gamma = -\kappa k/c_s$. However, it is useful to follow the connection between Cleary’s formula (70) and Eq. (63) obtained from the quasiclassical solution. In the classical limit the partial wave l corresponds to the quasiclassical trajectory with the impact parameter $b = l/k = \hbar l/p$ and the small scattering angle is $\varphi = -d\delta_l/dl = -(1/2\hbar k)d\delta S(b)/db$, where $\delta S(b)$ is the action variation along the trajectory with impact parameter b . Finally replacing the sum by an integral Eq. (70) can be rewritten as

$$\sigma_{\perp} = -\frac{2}{k} \int_{-\infty}^{\infty} dl \frac{d\delta_l}{dl} = -\frac{1}{\hbar k} \int_{-\infty}^{\infty} db \frac{d\delta S(b)}{db} . \quad (76)$$

which yields (63) since $\delta S_{\pm} = \delta S(\pm\infty)$. Strictly speaking Eq. (76) is valid if $\delta S(b)$ is a continuous function. This is the case for rotons which are scattered quasiclassically [5, 6, 7]. But it yields a correct answer even for phonons despite $\delta S(b)$ has a jump at $b \sim 0$. Thus even though phonon scattering cannot be described in the quasiclassical approximation (it yields the scattering angle $\varphi = -(1/\hbar k)d\delta S(b)/db \approx 0$ at $b \neq 0$), the quasiclassical expression (76) gives a correct phonon transverse cross-section.

8 Momentum balance in the two-fluid hydrodynamics

Up to now we analyzed spatial scales much less than the mean-free-path l_{ph} of phonons (ballistic region). Sound waves (phonons) interacted with the velocity field generated by a vortex, but phonon-phonon interaction was neglected. Now we shall see what is going on at scales much larger than l_{ph} , where the two-fluid hydrodynamics is valid.

Scattering of phonons by the vortex in the ballistic region produced a force $\vec{F}$ on the vortex, and according to the third Newton law a force $-\vec{F}$ on the phonons (the normal fluid) should also arise. A momentum transfer from a phonon to a vortex takes place at distances of the order of the phonon wavelength $\lambda = 2\pi/k$, which is much less than the mean-free path l_{ph} . This means that at hydrodynamic scales the force on the normal fluid is a δ -function force concentrated along the vortex line. The response of the normal fluid to this force is described by the Navier-Stokes equation with the dynamic viscosity η_n :

$$\frac{\partial \vec{v}_n}{\partial t} + (\vec{v}_n \cdot \vec{\nabla})\vec{v}_n = \nu_n \Delta \vec{v}_n - \frac{\vec{\nabla} P}{\rho} - \frac{\rho_s S}{\rho_n \rho} \vec{\nabla} T, \quad (77)$$

where $\nu_n = \eta_n/\rho_n$ is the kinematic viscosity, S is the entropy per unit volume, and T is the temperature. Our problem is similar to the Stokes problem of a cylinder moving through a viscous liquid [1]. Neglecting the nonlinear inertial (convection) term $(\vec{v}_n \cdot \vec{\nabla})\vec{v}_n$ in the Navier-Stokes equation, the δ -function force produces a divergent logarithmic velocity field (the Stokes paradox [1]):

$$\vec{v}_n(\vec{r}) = \vec{v}_n + \frac{\vec{F}}{4\pi\eta_n} \ln \frac{r}{l_{ph}}, \quad (78)$$

where $\vec{v}_n$ is the normal velocity at a distance of order $r \sim l_{ph}$, which separates the ballistic and the hydrodynamic regions. In this expression the mean-free path l_{ph} replaces the cylinder radius of the classical Stokes problem in the argument of the logarithm. Such a choice of the lower cut-off assumes that the quasiparticle flux on the vortex is entirely determined by the equilibrium Planck distribution at the border between the ballistic and the two-fluid-hydrodynamics region at $r \sim l_{ph}$ [37].

However small the relative normal velocity $\vec{v}_n - \vec{v}_L$ could be, at distance of the order or larger than $r_m \sim \nu_n/|\vec{v}_n - \vec{v}_L|$ the nonlinear convection term becomes important and stops a logarithmic growth of the normal velocity. Due to the force $\vec{F}$ the normal velocities $\vec{v}_{n\infty}$ and $\vec{v}_n$ at large ($r \sim r_m$) and small ($r \sim l_{ph}$) distances from the vortex line are different (the viscous drag [2]):

$$\vec{F} = \frac{4\pi\eta_n}{\ln(r_m/l_{ph})}(\vec{v}_{n\infty} - \vec{v}_n) . \quad (79)$$

At very large distances $r \gg r_m$ the nonlinear convection term is more important than the viscous term. Thus the scale r_m , which we shall call Oseen's length, divides the hydrodynamic region onto the viscous and convection subregions [38]. All relevant scales are shown in Fig 5. For a longitudinal force the solution of the Navier-Stokes equation valid for both the viscous and convection subregions was obtained by Oseen long ago (see Ref. [1]).

The force $-\vec{F}$ is transmitted to large distances by a constant momentum flux. In the viscous subregion momentum transport in the normal fluid occurs via viscosity: $F_i = -\oint \tau_{ij} dS_j$, where $\tau_{ij} = -\eta_n (\nabla_i v_{nj} + \nabla_j v_{ni})$ is the viscous stress tensor. On the other hand, the total momentum flux for the whole liquid should vanish: $\oint \Pi_{ij} dS_j + \oint \tau_{ij} dS_j = 0$, where Π_{ij} is given by Eq. (32). The normal velocity field does not contain the circular velocity $\vec{v}_v$, and there is no normal circulation in the viscous subregion. Therefore the momentum flux $\oint \Pi_{ij} dS_j$ yields the superfluid Magnus force, i.e., the momentum balance in the viscous subregion confirms that the force $\vec{F}$ satisfies Eq. (9).

In the convection region the viscosity becomes ineffective and the momentum conservation gives again $\oint \Pi_{ij} dS_j = 0$, like in the ballistic region. The superfluid part of the momentum flux is related to the superfluid Magnus force $\propto \rho_s$ and the normal part of the flux should transmit the same force $\vec{F}$. The relation between the force $\vec{F}$ and the relative normal velocity $\vec{v}_{n\infty} - \vec{v}_L$




Figure 5: Relevant scales: the phonon wavelength λ , the phonon mean free path l_{ph} , and Oseen's length $r_m \sim \nu_n/|\vec{v}_n - \vec{v}_L|$

in the convection subregion can be derived from Eqs. (10) and (79):

$$\vec{F} = -\tilde{D}(\vec{v}_L - \vec{v}_{n\infty}) - \tilde{D}'[\hat{z} \times (\vec{v}_L - \vec{v}_{n\infty})] , \quad (80)$$

where the parameters $\tilde{D}$ and $\tilde{D}'$ are connected with D and D' in Eq. (10) by a complex relation

$$\frac{1}{\tilde{D} + i\tilde{D}'} = \frac{\ln(r_m/l_{ph})}{4\pi\eta_n} + \frac{1}{D + iD'} . \quad (81)$$

Since in the convection region the viscosity is ineffective and the normal fluid behaves as an ideal one, the only way to transmit the transverse component of the force $\vec{F}$ to infinity is to create a circulation of the normal velocity [39]. Because of the viscous drag separation on the longitudinal and the transverse force should be done with respect to the normal velocity $\vec{v}_{n\infty} - \vec{v}_L$, but not $\vec{v}_n - \vec{v}_L$. Thus the normal circulation is determined by $\tilde{D}'$, but not D' :

$$\kappa_n = \oint d\vec{l} \cdot \vec{v}_n = -\frac{\tilde{D}'}{\rho_n} = \frac{\kappa}{1 + \left[\frac{\kappa\rho_n \ln(r_m/l_{ph})}{4\pi\eta_n} \right]^2} , \quad (82)$$

where we neglected the longitudinal component $\propto D$ in the ballistic region and used the relation $D' = -\kappa\rho_n$ for the Iordanskii force.

In the convection subregion more problematic is to transmit not the transverse, but the longitudinal component of the force. A longitudinal force is impossible if viscosity is neglected completely: a body moving through an ideal liquid does not produce any dissipative force (d'Alembert's paradox [40]). The paradox is resolved by finding that a *laminar wake* should arise behind a moving body. Within the wake one cannot neglect viscosity even deeply in the convection area when $r \gg r_m$. The width of the laminar wake is growing as $\sim \sqrt{rr_m}$ far from the moving body [1]. Solving the Navier-Stokes equation by Oseen's method one can find out how the laminar wake and the normal circulation are formed during the crossover from the viscous to the convection subregion. One can find a detailed analysis of this crossover in the presence of the longitudinal and the transverse force in Ref. [41].

The force $\vec{F}$ is the mutual friction force introduced by Hall and Vinen [2] for the analysis of propagation of the second sound in rotating superfluid. With help of Eqs. (9), (10), and (79) one obtains a linear relation between the mutual friction force and the counterflow velocity $\vec{v}_s - \vec{v}_{n\infty}$ bearing in

mind that the average normal velocity practically coincides with the velocity $\vec{v}_{n\infty}$:

$$\vec{F} = \frac{\kappa\rho_s\rho_n}{2\rho}B(\vec{v}_s - \vec{v}_{n\infty}) - \frac{\kappa\rho_s\rho_n}{2\rho}B'[\hat{z} \times (\vec{v}_s - \vec{v}_{n\infty})] , \quad (83)$$

The Hall-Vinen parameters B and B' are given by a complex relation [6]

$$\frac{2\rho}{\kappa\rho_s\rho_n}\frac{1}{B - iB'} = -\frac{1}{i\rho_s\kappa} + \frac{1}{\tilde{D} + i\tilde{D}'} = -\frac{1}{i\rho_s\kappa} + \frac{\ln(r_m/l_{ph})}{4\pi\eta_n} + \frac{1}{D + iD'} . \quad (84)$$

According to this relation a strong viscous drag (large logarithm $\ln(r_m/l)$, or small viscosity η_n) suppresses the effect of the transverse force ($\propto D'$) on the mutual friction [41]. But the effect of the superfluid Magnus force and the longitudinal force $\propto D$ is also suppressed in this limit. In fact, this is a limit when the force from quasiparticle scattering (transverse, or longitudinal) is so strong that the normal fluid sticks to the vortex, and $\vec{v}_L = \vec{v}_n$. Then the resultant force $\vec{F}$ depends only on viscosity, as in the classical Stokes problem.

9 Magnus force and the Berry phase

Now let us consider a connection between the transverse force and the Berry phase. We shall use the hydrodynamic description with the Lagrangian obtained from Eq. (13) after the Madelung transformation:

$$L = \frac{\kappa\rho}{2\pi}\frac{\partial\phi}{\partial t} - \frac{\kappa^2\rho}{8\pi}\nabla\phi^2 - \frac{V}{2}\rho^2 . \quad (85)$$

The first term with the first time derivative of the phase ϕ (Wess-Zumino term) is responsible for the Berry phase $\Theta = \Delta S_B/\hbar$, which is the variation of the phase of the quantum-mechanical wave function for an adiabatic motion of the vortex around a closed loop [16]. Here

$$\Delta S_B = \int d\vec{r} dt \frac{\kappa\rho}{2\pi}\frac{\partial\phi}{\partial t} = - \int d\vec{r} dt \frac{\kappa\rho}{2\pi}(\vec{v}_L \cdot \vec{\nabla}_L)\phi . \quad (86)$$

is the classical action variation around the loop and $\vec{\nabla}_L\phi$ is the gradient of the phase $\phi[\vec{r} - \vec{r}_L(t)]$ with respect to the vortex position vector $\vec{r}_L(t)$. However, $\vec{\nabla}_L\phi = -\vec{\nabla}\phi$, where $\vec{\nabla}\phi$ is the gradient with respect to $\vec{r}$. Then the loop

integral $\oint d\vec{l}$ yields the circulation of total current $\vec{j} = (\kappa/2\pi) < \rho \vec{\nabla} \phi >$ for points inside the loop, but vanishes for points outside. As a result, the Berry-phase action is given by [18]

$$\Delta S_B = V \frac{\kappa}{2\pi} \oint (d\vec{l} \cdot \vec{j}) . \quad (87)$$

where V is the volume inside the loop (a product of the loop area and the liquid height along a vortex). Contrary to Eq. (86), the loop integral in Eq. (87) is related with the variation of the position vector $\vec{r}$, the vortex position vector $\vec{r}_L$ being fixed.

Since the Berry phase is proportional to the current circulation, which determines the transverse force, there is a direct connection between the Berry phase and the amplitude of the transverse force on a vortex, as was shown, e.g., in Ref. [17]. Thus in order to find the Berry phase, one should calculate the current circulation. If the circulation of the normal velocity at large distances vanished (as assumed in Refs. [15, 17, 18]), the current circulation would be $\oint (d\vec{l} \cdot \vec{j}) = \rho_s \vec{\kappa}$, and the Berry phase (as well as the effective Magnus force) would be proportional to ρ_s . However, according to Sec. 8 and the recent analysis by Thouless *et al.* [41], at very large distances (in the convection subregion) the normal circulation $\kappa_n = \oint (d\vec{l} \cdot \vec{v}_n)$ does not vanish and $\oint (d\vec{l} \cdot \vec{j}) = \rho_s \vec{\kappa} + \rho_n \kappa_n$. Using a proper value of the asymptotic normal circulation κ_n given by Eq. (82) the Berry phase yields the same transverse force as determined from the momentum balance.

In order to obtain a correct value of the transverse force from the Berry phase, one should choose a loop radius much larger than Oseen's length r_m . If the loop radius is chosen in the viscous subregion $l_{ph} < r < r_m$, the total-current circulation is proportional ρ_s , but the Berry phase does not yield the total transverse force, since a part of it, namely, the Iordanskii force, is presented by the viscous momentum flux, which cannot be obtained in the Lagrange formalism. And if the loop radius is chosen in the ballistic region, the total-current circulation is not defined at all and depends on a shape of the loop, since the phonon mass current is not curl-free, as pointed out after Eq. (58).

10 Discussion and conclusions

The momentum-balance analysis definitely confirms an existence of the transverse force on a vortex from phonon scattering (the Iordanskii force). This conclusion agrees with the results of the recent analysis of Thouless *et al.* [41].

The Berry phase yields the same value of the transverse force as the momentum balance, if a proper value of the normal circulation at large distances from the vortex is used for the calculation of the Berry phase. However, the Berry-phase analysis itself cannot provide the normal-circulation value, since the latter is determined by the processes at small distances from the vortex, which are beyond of the Berry-phase analysis. The small-distance processes determine the force between the superfluid and the normal component, which is present in the small-distance boundary condition for the Navier-Stokes equation in the two-fluid-hydrodynamics region. The transverse force on the normal component is transmitted to infinite distances by the constant momentum flux, which requires a normal circulation far from the vortex. This circulation should be used for determination of the Berry phase.

Ambiguity of the Berry-phase analysis of the transverse force originates from ambiguity of the transverse force in the Lagrange formalism, which was discussed in the end of Sec. V in Ref. [7]). Adding a constant C to the total density in the Wess-Zumino term, i.e., replacing ρ by $\rho + C$, one obtain a different amplitudes of the Berry phase and the transverse force without any effect on the field equations for the condensate wave function (see also discussion of this constant in Fermi liquids by Volovik [42]). This arbitrariness has a profound physical meaning. Derivation of the Magnus force in the Lagrange formalism dealt only with large distances much exceeding the vortex core size. However, the processes inside the core affect the total Magnus force in general. Deriving the Magnus force from the momentum balance we use the condition that the total momentum flux $\oint \Pi_{ij} dS_j$ through the surface of large radius around the vortex line vanishes. This is true only for a Galilean invariant liquid satisfying the momentum conservation law. If the liquid in the vortex core interacts with the external world (e.g., with crystal impurities in superconductors), the momentum balance condition must be $\oint \Pi_{ij} dS_j = f_i$ where $\vec{f}$ is the force on the vortex core which could also have a transverse component (the Kopnin-Kravtsov force [14]). One can calculate such a force only from the vortex-core analysis. The results of this analysis

must be used for determination of an unknown constant in the Wess-Zumino term.

Acknowledgements

I thank N. Kopnin, L. Pitaevskii, A. Shelankov, M. Stone, D. Thouless, and G. Volovik for interesting discussions. The work was supported by the grant of the Israel Academy of Sciences and Humanities.

References

- [1] H. Lamb, *Hydrodynamics* (Cambridge University Press, New York 1975).
- [2] H.E. Hall and W.F. Vinen, Proc. Roy. Soc. **A238**, 204 (1956).
- [3] E.M. Lifshitz and L.P. Pitaevskii, Zh. Eksp. Teor. Fiz. **33**, 535 (1957) [Sov. Phys.-JETP **6**, 418 (1958)].
- [4] S.V. Iordanskii, Zh. Eksp. Teor. Fiz. **49**, 225 (1965) [Sov. Phys.-JETP **22**, 160 (1966)].
- [5] E.B. Sonin, Zh. Eksp. Teor. Fiz. **69**, 921 (1975) [Sov. Phys.-JETP **42**, 469 (1976)].
- [6] E.B. Sonin, Rev. Mod. Phys. **59**, 87 (1987).
- [7] E.B. Sonin, Phys. Rev. B **55**, 485 (1997).
- [8] Y. Aharonov and D. Bohm, Phys. Rev. **115**, 485 (1959).
- [9] M.V. Berry, R.G. Chambers, M.D. Large, C. Upstill, and J.C. Walmsley, Eur. J. Phys. **1**, 154 (1980).
- [10] P. Roux, J. de Rosny, M. Tanter, and M. Fink, Phys. Rev. Lett. **79**, 3170 (1997).
- [11] U. Leonhardt and P. Piwnicki, Phys. Rev. Lett. **84**, 822 (2000).
- [12] A.L. Shelankov, this book.

- [13] P. Nozières and W.F. Vinen, *Phil. Mag.* **14**, 667 (1966).
- [14] N.B. Kopnin, this book.
- [15] P. Ao and D.J. Thouless, *Phys. Rev. Lett.* **70**, 2158 (1993).
- [16] M.V. Berry, *Proc. R. Soc. London A* **392**, 45 (1984).
- [17] D.J. Thouless, P. Ao, and Q. Niu, *Phys. Rev. Lett.* **76**, 3758 (1996).
- [18] M.R. Geller, C. Wexler, and D.J. Thouless, *Phys. Rev. B* **57**, R8119 (1998).
- [19] H.E. Hall and J.R. Hook, *Phys. Rev. Lett.*, **80**, 4356 (1998); C. Wexler, D.J. Thouless, P. Ao, and Q. Niu, *ibid.*, **80**, 4357 (1998).
- [20] E.B. Sonin, *Phys. Rev. Lett.* **81**, 4276 (1998).
- [21] C. Wexler, D.J. Thouless, P. Ao, and Q. Niu, *Phys. Rev. Lett.* **81**, 4277 (1998).
- [22] A short version of this paper is to be published in *Proceedings of the Isaac Newton Institute Workshop on Quantized Vortex Dynamics and Superfluid Turbulence*, Cambridge, UK, August 2000.
- [23] E.P. Gross, *Nuovo Cimento* **20**, 454 (1961); L.P. Pitaevskii, *Zh. Eksp. Teor. Fiz.* **40**, 646 (1961) [*Sov. Phys.-JETP* **13**, 451 (1961)].
- [24] But this assumption becomes invalid for the roton normal fluid at higher temperatures, especially close to the critical temperature. Then other methods based on the modified Ginzburg-Pitaevskii theory should be used [6, 25].
- [25] E.B. Sonin, *J. Low Temp. Phys.* **42**, 417 (1981).
- [26] N.B. Kopnin, *Physica B* **210**, 267 (1995).
- [27] T.D.C. Bevan *et al*, *J. Low Temp. Phys.* **109**, 423 (1997).
- [28] R.J. Donnelly, *Quantized vortices in helium II* (Cambridge University Press, Cambridge, 1991), Sec. 2.8.3.

- [29] R. Peierls, *Surprises in Theoretical Physics*, (Princeton, 1979).
- [30] M.Stone, Phys. Rev. E **62**, 1341 (2000).
- [31] S.J. Putterman and P.H. Roberts, Physica **117** A, 369 (1983).
- [32] L.P. Pitaevskii Zh. Eksp. Teor. Fiz. **35**, 1271 (1958) [Sov. Phys.-JETP **8**, 888 (1959)].
- [33] A. Fetter, Phys. Rev. **136A**, 1488 (1964).
- [34] A.L. Shelankov, Europhys. Lett., **43**, 623 (1998)
- [35] M. Stone, Phys. Rev. B **61**, 11 780 (2000).
- [36] R.M. Cleary, Phys. Rev. **175**, 587 (1968).
- [37] However, scattered quasiparticles colliding with other quasiparticles immediately after scattering can return back and modify the quasiparticle flux on the vortex. As a result of it, the lower cut-off could be a combination of the mean-free path l_{ph} and some effective cross-section of the vortex. This logarithmically weak effect was shortly discussed in the end of Sec. 1 in Ref. [5].
- [38] Note that the choice of Oseen's length as a upper cut-off of the logarithmic factor is valid for an isolated vortex under a stationary fluid flow. In general the upper cut-off should be the smallest among three scales: Oseen's length, the frequency-dependent viscous depth, and the intervortex spacing [6].
- [39] Existence of the normal circulation at large distances in the presence of the transverse force on the vortex was pointed out by Pitaevskii (unpublished).
- [40] L.D. Landau and E.M. Lifshitz, *Fluid Mechanics*, (Pergamon Press, Oxford 1982).
- [41] D.J. Thouless, M.R. Geller, W.F. Vinen, J-Y. Fortin, and S.W. Rhee, cond-mat/0101297.
- [42] G.E. Volovik, Pis'ma v ZhETF, **65**, 641 (1997) [JETP Letters, **65**, 676 (1997)].